

---

# OPHTHALMOLOGICAL SOCIETY OF THE WEST INDIES

**Proceedings of the  
24<sup>th</sup> Annual Congress  
July 10 – 13, 2013  
Royal Torarica Hotel  
Paramaribo, Suriname**



**Editor-in-Chief**  
*Professor Everard Barton*

## **Congress Committee and Scientific Editors**

*Dr Lizette Mowatt  
Dr Sonja Johnston  
Dr Desiree Murray  
Dr Hugh Vaughn  
Dr Jeffery J Hurwitz*

*Dr Angelo Ramirez  
Dr Donovan Calder  
Dr Alfred L Anduze  
Dr DC Gibbons  
Dr A Bueno de Mesquita-Voigt*

## **OSWI COUNCIL 2012 – 2014**

<i>President</i>	–	<i>Dr Lizette Mowatt</i>
<i>Vice-President</i>	–	<i>Dr Angelo Ramirez</i>
<i>Secretary/Treasurer</i>	–	<i>Dr Sonja Johnston</i>
<i>Immediate Past President</i>	–	<i>Dr Terrence Allan</i>
<i>Assistant Secretary</i>	–	<i>Dr Alfred Anduze</i>
<i>Immediate Past President</i>	–	<i>Dr Donovan Calder</i>



## OUR PLASMA SURFACE DEFENDS AGAINST DAILY DEPOSITS.

All **AIR OPTIX®** brand contact lenses have a unique surface technology that's proven to maintain wettability<sup>1\*</sup> and resist deposits better than other available two-week or monthly replacement SiHy lenses.<sup>2,3\*\*</sup>

Superior Surface With Moisture & Consistent Comfort



## AIR OPTIX® BRAND Family of Contact Lenses

See our superior surface deposits resistance and wettability<sup>1-3\*\*</sup> data at [MYALCON.COM](http://MYALCON.COM)



<sup>1</sup>Compared to ACUVUE 1 CONTACT, ACUVUE 2 ADVANCE, ACUVUE 2 HYDRONIGHT, ACUVUE 2 WETBOND, ACUVUE 2 and ACUVUE 2 OASYS contact lenses. <sup>2</sup>Superior lipid deposit resistance compared to ACUVUE 1 CONTACT, ACUVUE 2 ADVANCE, ACUVUE 2 HYDRONIGHT, ACUVUE 2 WETBOND, ACUVUE 2 OASYS contact lenses. <sup>3</sup>Superior protein deposit resistance compared to ACUVUE 1 CONTACT, ACUVUE 2 ADVANCE, ACUVUE 2 HYDRONIGHT, ACUVUE 2 WETBOND, ACUVUE 2 OASYS contact lenses. \*Wettability for illustrative purposes and not an exact representation. \*\*Trademarks are the property of their respective owners.

**Important information for AIR OPTIX® AQUA (Jotraficon B), AIR OPTIX® AQUA Multifocal (Jotraficon B) and AIR OPTIX® for Astigmatism (Jotraficon B) contact lenses:** Do not wear or extended wear up to 6 nights for new/long-time wearers, prodrops and/or astigmatism. Risk of serious eye problems (e.g., corneal ulcer) is greater for extended wear. In rare cases, loss of vision may result. Side effects like discomfort, mild burning or stinging may occur.

**Important information for AIR OPTIX® NIGHT & DAY® AQUA (Jotraficon A) contact lenses:** Intended for vision correction for daily wear (wear only while awake) or extended wear (wear while awake and asleep) for up to 30 nights. **Relevant Warnings:** A corneal ulcer may develop rapidly and cause eye pain, redness or blurry vision with progressive, light sensitivity, usual, and/or increased loss of vision, may result. The risk of serious problems is greater for extended wear vs. daily wear and smoking increases this risk. A one-year post-market study found 0.10% (10 out of 10,000) of wearers developed a severe corneal infection, with 0.04% (4 out of 10,000) of wearers experiencing a permanent reduction in vision by two or more rows of letters on an eye chart.

**Relevant Precautions:** Not every eye can wear for 30 nights. Approximately 80% of wearers can wear the lenses for extended wear. About two-thirds of wearers achieve the full 30-night continuous wear. **Side Effects:** In clinical trials, approximately 3-5% of wearers experienced at least one episode of irritative keratitis, a localized inflammation of the cornea which may be accompanied by mild to severe pain and may require the use of antibiotic eye drops for up to one week. Other less serious side effects were conjunctivitis, lid irritation or lens discomfort including dryness, mild burning or stinging. **Contraindications:** Contact lenses should not be worn if you have eye infection or inflammation (redness and/or swelling), eye disease, injury or dryness that interferes with contact lens wear; eye disease (e.g., that may be affected by or impact lens wear); certain allergic conditions or using certain medications (e.g., certain eye medications). **Additional information:** Lenses should be replaced every month. If removed before then, lenses should be cleaned and disinfected before wearing again. Always follow the eye care professional's recommended lens wear, care and replacement schedule. Consult package insert for complete information, available without charge by calling (800) 241-3999 or go to [myalcon.com](http://myalcon.com).

**References:** 1. In vitro measurement of contact lens surface wettability; significance demonstrated at the 0.001 level; Alcon data on file, 2009. 2. Nash W, Gabriel M, Murray M, Koo M. A comparison of various silicone hydrogel lenses: lipid and protein deposition as a result of daily wear. Optom Vis Sci. 2010;87:E abstract 103110. 3. In vivo measurement of lipid deposits on lenses worn daily wear through consecutive measurement of replacement periods; (C) AIR OPTIX® 1. Cleaning and disinfecting solution used for cleaning and disinfecting; significance demonstrated at the 0.05 level; Alcon data on file, 2008.

See product instructions for complete wear, care, and safety information.

## CONTENTS

---

---

Acknowledgements .....	1
Preface .....	2
Programme .....	3
Abstracts	
Session 1 – Workshop on Oculoplastics .....	9
Session 2 – Workshop on Lifestyle Management .....	10
Session 3 – Advocacy Issues/Public Health .....	12
Opening Ceremony	
Session 4 – Dr Garth Taylor Memorial Lecture .....	13
Session 5 – Medical Retina .....	14
Session 6 – Surgical Retina .....	16
Session 7 – Glaucoma .....	17
Session 8 – Cataract/Refractive Surgery .....	19
Session 9 – Cornea and External Eye Diseases .....	21
Session 10 – Residents' Presentations .....	23
Session 11 – Live Surgeries – Suriname Eye Centre .....	26
Poster Presentations .....	27
Ophthalmic Assistants, Nurses and Technicians Programme .....	29
Ophthalmic Assistants, Nurses and Technicians Abstracts .....	33
Index of Participants .....	35