

A Hantavirus Case with Confusion and Diffuse Conjunctival Haemorrhage

The Editor,

Sir,

A 19-year-old male carpenter living in a village affiliated to the district of Görele in the province of Giresun, Turkey, presented to the hospital with symptoms of fever and confusion. Cranial CT and cerebrospinal fluid's findings were normal. ARB and Herpes PCR tests from CSF were negative, and there was no growth in CSF. No characteristic was identified at complete blood count apart from thrombocytopenia ($82 \times 10^3/\mu\text{L}$). A preliminary diagnosis of acute meningitis was made, and the patient was started on acyclovir and ceftriaxone therapy. His confusion resolved in 24 h. Elevation was determined in urea and creatinine, but there was no growth in blood cultures. Thrombocyte values decreased to $32 \times 10^3/\mu\text{L}$ at third day. Bilateral retinal hemorrhage and kidney function impairment developed.

At laboratory analysis, complete blood count was WC: $8300 \times 10^3/\mu\text{L}$, hemoglobin: 11.9 g/dL, thrombocyte: $142 \times 10^3/\mu\text{L}$ and sedimentation rate: 34 mm/s (0–20). Serum CRP was 4.4 mg/dL (< 0.5), procalcitonin: 0.31 $\mu\text{g/L}$ (< 0.5), PT: 13.2 sec (11–15.5), aPTT: 32.8 sec (23.6–34.8), INR: 1.07 (0.8–1.25) BUN: 21 mg/dL (0–21), creatinine: 1.74 mg/dL (0.67–1.17), AST: 60 U/L (0–35), ALT: 24 U/L (0–45), total bilirubin: 0.6 mg/dL (0.3–1.2), LDH: 561 U/L (<248), CK: 311 U/L (20–200), myoglobin: 67.75 ng/mL (28–72), total protein: 5.8 g/dL (6.6–8.3) and albumin: 2.8 g/dL (3.5–5.2). Complete urine examination was normal. The brucella agglutination test was negative. The MAT test performed for leptospirosis was also negative.

Further investigation of the serum samples at the Virology Reference and Research Laboratory, Public Health Institute of Turkey by IFA (Anti-hantavirus IIFT, LIFT, Germany) and the results were confirmed by immunoblot (Hantavirus Profile 1 Euroline IgM, IgG Euroimmun, Germany). The patient was diagnosed hemorrhagic fever with renal syndrome (HFRS).

Since the Dobrava serotype carried by the yellow-necked wood mouse (*Apodemus flavicollis*) is able to grow in moist soil at low temperatures and levels of ultraviolet light, cases from Turkey have largely been reported from the Black Sea region. The first cases diagnosed with HFRS from the area of Giresun were reported by us in 2010, and similar case reports have followed in subsequent years (1, 2). Complications involving the eyes and central nervous system have previously been reported in some cases of HFRS (3, 4). Since confusion had entirely resolved after time of admission to our department, CSF examination was not required in this case. CT of the brain was normal. Since fever, conjunctival hemorrhage and polyuria resolved entirely at follow-up and laboratory values were normal, the patient was discharged in a healthy condition on the 10th day

In conclusion, it should keep in mind that hantaviruses may also lead to clouding of consciousness without intracranial hemorrhage and those infections which develop with Dobrava virus may have more serious clinical findings comparing to other subtypes.

AUTHORS' NOTE

Authors declare that there is no actual or potential conflict of interest in relation to this article.

Keywords: Confusion, Dobrava-Belgrade virus, haemorrhage

S Kaya¹, AM Şahin², DY Çağlayık³, G Yılmaz¹, İ Köksal¹

From: ¹Department of Infectious Diseases and Clinical Microbiology, School of Medicine, Karadeniz Technical University, Trabzon, ²Giresun State Hospital, Infectious Diseases Clinic, Giresun, and ³Department of Microbiology Reference Laboratories, Virology Reference and Research Laboratory, Public Health Institute of Turkey, Ankara, Turkey.

Correspondence: Dr. S Kaya, Department of Infectious Diseases and Clinical Microbiology, School of Medicine, Karadeniz Technical University, Trabzon, Turkey. Fax: +904623775344, e-mail: eselkaya@gmail.com

REFERENCES

1. Reusken C, Heyman P. Factors driving hantavirus emergence in Europe. *Curr Opin Virol* 2013; **3**: 92–9.
2. Kaya S, Yılmaz G, Erensoy Ş, Çağlayık DY, Uyar Y, Köksal İ. Doğu Karadeniz, Giresun İlinde tespit edilen iki hantavirüs enfeksiyonu olgusu. *Mikrobiyol Bul* 2010; **44**: 479–87.
3. Hautala T, Hautala N, Mähönen SM, Sironen T, Pääkkö E, Karttunen A et al. Young male patients are at elevated risk of developing serious central nervous system complications during acute Puumala hantavirus infection. *BMC Infect Dis* 2011 Aug 14; **11**: 217. doi: 10.1186/1471-2334-11-217.
4. Hautala N, Kauma H, Rajaniemi SM, Sironen T, Vapalahti O, Pääkkö E et al. Signs of general inflammation and kidney function are associated with the ocular features of acute Puumala hantavirus infection. *Scand J Infect Dis* 2012; **44**: 956–62. doi: 10.3109/00365548.2012.700119. Epub 2012 Jul 25.