

## Breast Cancer Receptor Profiles in Jamaica: A 6-year Analysis

SN Chin<sup>1</sup>, CA Walters<sup>2</sup>, E Williams<sup>1</sup>

### ABSTRACT

**Objective:** To determine the breast cancer immunohistochemistry (IHC) receptor status for tests performed at the University of the West Indies (UWI) from January 2002 to December 2007 and to investigate for an association between receptor profile and patient age, tumour grade and stage.

**Methods:** The UWI breast cancer IHC receptor database was examined to determine receptor profile, patient age, tumour histology, grade, size and lymph node status.

**Results:** In total, 1383 breast cancer cases were tested for oestrogen receptor (ER) and human epidermal growth factor receptor 2 (HER 2) statuses during the study period; progesterone receptor (PR) testing was not available. Receptor profiles were: ER+/HER2- (50.2%), ER-/HER2- (28.1%), ER+/HER2+ (15.3%) and ER-/HER2+ (6.4%). Across all age groups, ER+/HER2- was the most frequent profile (45–52%) and ER-/HER2- was the second most frequent (27–34%). There was no statistically significant association between the receptor profile and age ( $p = 0.079$ ). Amongst Grade III tumours, ER-/HER2- was the most prevalent profile (44.6%); ER+/HER2- was the most prevalent for Grade I and Grade II tumours (60.7 and 48.8%, respectively). There was a statistically significant association between the receptor profile and tumour grade ( $p = < 0.001$ ). There was no statistically significant association between the receptor profile and tumour stage ( $p = 0.359$ ).

**Conclusion:** The prevalence of ER/HER2-negative breast cancer was 28%, in keeping with triple-negative breast cancer (TNBC) prevalence in African-American populations. There was a statistically significant association between the receptor profile and tumour grade ( $p < 0.001$ ) (most Grade III tumours were ER-/HER2-), in keeping with the biologically aggressive behaviour of TNBC.

**Keywords:** Breast cancer receptors, immunohistochemistry, Jamaica.

### INTRODUCTION

Breast cancer is the most common malignancy among Jamaican women, with an age-standardized incidence rate of 43.1 per 100 000 (1). Breast cancer prognostication has traditionally been derived from tumour-related histopathological variables of size, histological grade and nodal status. The Nottingham Prognostic Index is a well-validated prognostic system that uses these criteria to determine prognosis following surgery for breast cancer (2). While these pathological criteria remain relevant, breast cancer molecular profiling has become increasingly important in prognostication. Biomarkers

such as the oestrogen receptor (ER), progesterone receptor (PR) and human epidermal growth factor receptor 2 (Her2)-neu oncogene (HER2) are not only important prognostic markers but are also predictive of response to targeted therapies (3). The triad of ER, PR and HER2 status has therefore become a mandatory part of therapeutic decision making in breast cancer management (4), with the predictive value of these biomarkers influencing the choice of systemic therapy.

In Jamaica, breast cancer receptor status testing is centralized to the Immunohistochemistry (IHC) Laboratory in the Department of Pathology, University

From: <sup>1</sup>Department of Pathology, University of the West Indies, Mona, Jamaica and <sup>2</sup>Office of the Dean, Faculty of Medical Sciences, University of the West Indies, Mona, Jamaica.

Correspondence: Dr SN Chin, Department of Medicine, University Hospital of the West Indies, Kingston 7, Jamaica.  
Email: sheray.chin@uwimona.edu.jm

of the West Indies (UWI), which undertakes testing for the entire island. The laboratory receives requests for IHC breast receptor studies from the University Hospital of the West Indies (UHWI), the teaching hospital with which the UWI is affiliated, the National Public Health laboratory (the government laboratory facility), regional pathology laboratories (eg, Mandeville Regional Hospital, Cornwall Regional Hospital) and several private pathology laboratories.

The UWI Department of Pathology IHC breast receptor database was established in 2002 and records results of tests performed on tumour tissue from patients with breast cancer. In addition, the database contains records from the histopathology report for each patient, including age, tumour histological type, grade, size and lymph node status (if nodal sampling was performed).

In this report, we will present results from the breast cancer IHC breast receptor database for tests performed during the 6-year period following its establishment. During this period, breast IHC breast receptor testing was limited to ER and HER2 (PR testing was not offered); hence, only results of ER and HER2 are available and will be presented.

## SUBJECTS AND METHODS

We reviewed the results of all IHC studies for the assessment of breast cancer expression of ER and HER2, performed at the UWI between January 2002 and December 2007. The biopsy specimens were from patients diagnosed at the UHWI as well as cases referred from other hospitals. The samples included breast core biopsies and mastectomy specimens. The IHC breast receptor database was examined to determine the ER/HER2 receptor profiles for all breast cancer specimens submitted for receptor testing during the study period, to record patient age, tumour histological type and grade, as well as tumour size and lymph node status if nodal sampling was performed.

### Protocol for immunostaining for oestrogen receptor and HER-2-Neu

The fixative for all internal and external cases was 10% phosphate-buffered formalin. The most appropriate block for performance of IHC studies was chosen after review of the H&E-stained slides. Paraffin sections were cut at 2–4  $\mu$ m, mounted on slides coated with poly-L-lysine solution, dried and then deparaffinized in xylene and rehydrated in decreasing strengths of ethanol to deionized water, then rinsed in phosphate-buffered saline. Blocking of non-specific background binding

was carried out using Universal Blocking Reagent. Incubations were performed with primary monoclonal antibodies: heat shock protein (HSP) 27 (Biogenex, Cat. No. AM171-5M) and HER-2-neu (Biogenex, Cat. No. AM 134-5M). Antigen/Antibody interaction was developed using anti-mouse immunoglobulin and conjugated streptavidin as a substrate for HSP and diaminobenzidine for HER-2/neu. The primary monoclonal antibody was omitted in the negative control and replaced by phosphate-buffered saline.

## Scoring system

### *Oestrogen receptor*

Positivity for ER was scored on a scale of I–III based on the number of cells with bright red cytoplasmic staining as follows: Grade I—few positive cells, Grade II—moderate number of positive cells and Grade III—many positive cells. The absence of any cell staining was interpreted as a negative result.

### *HER 2-Neu*

Human epidermal growth factor receptor -2 receptor status was tested with the HercepTest<sup>®</sup>, with scores reported on a scale of 0–3+ as follows: Score 0 (no membrane staining, or less than 10%), 1+ (faint membrane staining in more than 10% of tumour cells), 2+ (weak to moderate complete membrane staining in more than 10% of tumour cells) and 3+ (strong complete membrane staining in more than 10% of tumour cells) (5). Human epidermal growth factor receptor 2 protein overexpression was assessed as negative for a score of 0 or 1+, and positive for a score of 3+. Scores of 2+ required confirmatory testing with fluorescent in situ hybridization (FISH) for final determination. A positive result was defined as more than six HER2 gene copies per nucleus or a FISH ratio (HER2 gene signals to chromosome 17 signals) of more than 2.2; a negative result was a FISH result of less than four HER2 gene copies per nucleus, or FISH ratio of less than 1.8 (6).

Univariate analyses with descriptive summary statistics are presented for relevant variables. Chi-square tests for association between different receptor profiles and patient age, tumour grade and tumour stage were done using Stata version 13.0. The UHWI/UWI/Faculty of Medical Sciences Ethics Committee approved this study.

## RESULTS

Immunohistochemical studies for the assessment of expression of ER and HER2 were performed on 1383

breast cancer cases during the period 2002–2007. Table 1 summarizes the clinicopathological data for these patients. Median age was 52 years (range 20–94 years, IQR 43–63). Immunohistochemistry receptor profiles were: ER+/HER2– (50.2%), ER–/HER2– (28.1%), ER+/HER2+ (15.3%) and ER–/HER2+ (6.4%) (Table 2). Histopathological staging details, including tumour size and nodal status, were available for 860 patients and are presented in Table 3. The majority of tumours were greater than 2 cm in size (T2 or greater), and 67% had lymph node metastases. The aggregate pTNM stage was derived for the 439 cases with both tumour size and nodal status available and was: Stage I (5%), Stage II (50%) and Stage III (45%).

Table 1: Clinicopathological characteristics of patients with breast cancer IHC testing performed 2002–2007

	Total (n = 1383)
<b>Age</b>	
Range	20–94
Median (IQR)	52 (43–63)
<b>Age group</b>	
40 and under	240 (17.3%)
41–60	730 (52.8%)
61–70	214 (15.5%)
71 and over	199 (14.4%)
<b>Sex</b>	
Female	1377 (99.6%)
Male	6 (0.4%)
<b>Specimen type</b>	
Biopsy	708 (51.2%)
Mastectomy	675 (48.8%)
<b>Histology</b>	
In situ ductal	110 (8.0%)
In situ lobular	4 (0.3%)
Infiltrating ductal	1148 (83.0%)
Infiltrating lobular	108 (7.8%)
Mixed ductal/lobular	13 (0.9%)
<b>Tumour grade (BRS)</b>	
I	208 (15.1%)
II	486 (35.1%)
III	257 (18.6%)
Not graded	432 (31.2%)

IHC = immunohistochemistry; IQR = interquartile range.

Correlative analysis results are presented in Tables 4–6. Across all age groups, ER+/HER2– was the most frequently seen profile (45%–52%), with the ER–/HER2– profile being the second most frequently seen (27%–34%). There was no statistically significant association between the receptor profile and age (Chi-squared test statistic = 15.4453;  $df = 9$ ;  $p$ -value = 0.079).

Table 2: Receptor profiles results for breast cancer IHC testing during period 2002–2007

Receptor profile	No of patients (%)
ER+/Her2+	212 (15.3%)
ER+/Her2-	694 (50.2%)
ER–/Her2+	88 (6.4%)
ER–/Her2-	389 (28.1%)
<b>Total</b>	1383

IHC = immunohistochemistry; ER = oestrogen receptor; HER2 = human epidermal growth factor receptor 2.

Table 3: Histopathological staging for breast cancer cases with IHC testing (2002–2007)

	n = 860
<b>Tumour size, cm</b>	
Range	0–15
Median (IQR)	3 (2–5)
<b>Lymph nodes examined, no</b>	
Range	0–22
Median (IQR)	4 (0–10)
<b>Tumour size (TNM)</b>	
T0	44 (5.1%)
T1	162 (18.8%)
T2	444 (51.6%)
T3	210 (24.4%)
<b>Nodal status (TNM) (n = 458)</b>	
N0	154 (33.6%)
N1	150 (32.8%)
N2	112 (24.5%)
N3	42 (9.2%)
<b>TNM stage (n = 439)</b>	
I	22 (5.0%)
IIA	111 (25.3%)
IIB	110 (25.1%)
IIIA	154 (35.1%)
IIIC	42 (9.6%)

IHC = immunohistochemistry; IQR = interquartile range.

Table 4: Receptor profile according to patient's age

	No of patients within each age range (%)			
Receptor profile	40 and under	41–60	61–70	71 and over
ER+/Her2+	23 (17.0%)	70 (15.9%)	14 (9.4%)	16 (11.7%)
ER+/Her2-	61 (45.2%)	202 (46.0%)	77 (51.7%)	61 (44.5%)
ER–/Her2+	14 (10.4%)	35 (8.0%)	8 (5.4%)	5 (3.7%)
ER–/Her2–	37 (27.4%)	132 (30.1%)	50 (33.6%)	55 (40.1%)
<b>Total</b>	135	439	149	137

Chi-squared test statistic = 15.4453;  $df = 9$ ;  $p$ -value = 0.079; ER = oestrogen receptor; HER2 = human epidermal growth factor receptor 2.

Tumour grade was recorded for 658 cases. Amongst Grade III tumours, the most commonly seen receptor profile was ER–/HER2– (44.6%). For both Grade I

Table 5: Receptor profile according to tumour's grade

Receptor profile	No of patients within each tumour grade (%)		
	I	II	III
ER+/Her2+	13 (8.4%)	52 (15.9%)	19 (10.9%)
ER+/Her2-	94 (60.7%)	160 (48.8%)	65 (37.1%)
ER-/Her2+	3 (1.9%)	27 (8.2%)	13 (7.4%)
ER-/Her2-	45 (29.0%)	89 (27.1%)	78 (44.6%)
<b>TOTAL</b>	<b>155</b>	<b>328</b>	<b>175</b>

Chi-squared test statistic = 32.7394;  $df = 6$ ;  $p$ -value < 0.001 (excluding 'Not Graded'); ER = oestrogen receptor; HER2 = human epidermal growth factor receptor 2.

Table 6: Receptor profile according to tumour's stage

Receptor profile	No of patients within each tumour stage (%)		
	I	II	III
ER+/Her2+	2 (9.1%)	22 (9.9%)	31 (15.8%)
ER+/Her2-	11 (50.0%)	101 (45.7%)	77 (39.3%)
ER-/Her2+	0 (—)	17 (7.7%)	18 (9.2%)
ER-/Her2-	9 (40.9%)	81 (36.7%)	70 (35.7%)
<b>Total</b>	<b>22</b>	<b>221</b>	<b>196</b>

Chi-squared test statistic = 6.6; ER = oestrogen receptor; HER2 = human epidermal growth factor receptor 2.

and Grade II tumours, the ER+/Her2- profile was most common (60.7 and 48.8%, respectively). There was a statistically significant association between the receptor profile and tumour grade (Chi-squared test statistic = 32.7394;  $df = 6$ ;  $p$ -value < 0.001).

The majority of ER-/HER2- tumours were Stage II (51%) or Stage III (44%). Most ER+/HER2- tumours were also Stage II (53%) or Stage III (41%). There was no statistically significant association between the receptor profile and tumour stage (Chi-squared test statistic = 6.6078;  $df = 6$ ;  $p$ -value = 0.359).

## DISCUSSION

This 6-year analysis of breast receptor profiles from our IHC database represents the first study of the IHC receptor profiles of breast cancer from multiple centres across Jamaica. Previous studies of patients with breast cancer treated at the UHWI (7, 8) have demonstrated rates similar to that which we found in this larger multi-centre study group, in which cases from the entire island underwent testing at a centralized IHC testing facility. The rate of ER positivity in our study was 66%, compared to 62% (7) and 63% (8) in previous reports, while 22% of cases were HER2 positive, similar to previous reports of 20% HER2 positivity in patients treated at the UHWI (7).

The triple-negative phenotype (ER, PR and HER2 negative) is a surrogate for basal-like breast cancer and is known to be more aggressive and associated with a

relatively poorer prognosis (9). In population-based studies, basal-like breast cancer has been shown to have a higher prevalence in African American compared with non-African American patients (10), comprising 26 and 16% of cases in African American and non-African American women, respectively. Stead *et al* showed a 3-fold higher risk of triple-negative breast cancer (TNBC) (95% CI 1.6, 5.5;  $p = 0.0001$ ) in black compared with white women. In their study, the overall rate of TNBC was 20%; 30% of tumours in black women were triple negative, compared with 11%–13% of tumours in other women (11). There have been few studies examining the hormone receptor profile in African patients. In a study of 120 cases out of Kenya, Bird *et al* noted that 24% had ER- positive tumours, 34% were ER-negative and/or PR-positive, 10% were ER-negative but PR-positive and 66% were negative for ER and PR (12).

The Jamaican population is diverse in ethnicity, with a strong Afro-centric background. Based on this, we postulated that the rates of TNBC may be relatively high in our population. As previously noted, PR testing was not done during the study period, limiting our ability to comment on the prevalence of TNBC. Despite this limitation, we found a prevalence of 28% for ER/HER2-negative disease, which is in keeping with TNBC prevalence in the aforementioned North American studies. Despite the African influence on Jamaican ethnic background, it is interesting that our breast cancer hormone expression profile is more similar to that seen in African Americans in North American studies.

Since the period under study, PR testing has been implemented, in keeping with guideline recommendations that determination of tumour ER/PR and HER2 status should be done on all breast cancers (13). As we continue to add to our database with more complete information on PR, we will re-visit these prevalence rates.

We were unable to show a statistically significant association between receptor profile and patient age, which was anticipated based on the known propensity of basal like and TNBC for younger age groups (10). Aggressive tumour features such as high histological grade are characteristic of TNBC, and our study showed a significant association between receptor profile and tumour grade ( $p$ -value < 0.001).

Human epidermal growth factor receptor 2-overexpressing breast cancer, accounting for 22% of cases in this study, is a disease-subset associated with more aggressive behaviour, with significantly shortened disease-free survival and overall survival (14). Recognition of the

poor prognosis of HER2-positive breast cancer (15) has driven research and development of several anti-HER2 agents. Trastuzumab, a monoclonal antibody targeting the extracellular domain of HER2 protein, was shown to increase the clinical benefit of first-line chemotherapy in metastatic HER2 over-expressing breast cancer (16). This drug was approved by the The United States Food and Drug Administration for the adjuvant treatment of HER2-positive breast cancer based on the significant increases in both disease free and overall survival with its addition to standard adjuvant chemotherapy. In the combined analysis of two landmark Phase III trials, NSABP B-31 and the NCCTG N9831, adding trastuzumab to chemotherapy resulted in a significant improvement in DFS (HR: 0.48; 95% CI: 0.39, 0.59;  $p < 0.0001$ ) and a 33% reduction in the risk of death ( $p = 0.015$ ) (17). Based on our data, trastuzumab therapy may be indicated for up to 22% of our breast cancer patients. In our experience, however, only a limited number of eligible patients are able to receive this treatment primarily due to high drug cost. With such a potential gain in survival, a cost-benefit analysis is encouraged to support implementation of a trastuzumab subsidy by the national insurance scheme (National Health Fund).

## CONCLUSION

The prevalence of ER-negative/HER2-negative breast cancer in Jamaica (28%) is in keeping with TNBC prevalence in African American populations in North American studies. Inclusion of PR testing as part of breast IHC receptor testing will help us better define the prevalence of TNBC, and treatment strategies aimed at this aggressive subset of breast cancer must be pursued locally. There was a statistically significant association between the receptor profile and tumour grade ( $p = < 0.001$ ). Over 20% of breast cancers studies over-expressed HER2, and efforts aimed at increasing accessibility of targeted anti-HER2 therapy are encouraged.

## REFERENCES

1. TN Gibson, B Hanchard, N Waugh, D McNaughton. Age-specific incidence of cancer in Kingston and St Andrew, Jamaica, 2003–2007. *West Indian Med J* 2010; **59**: 456–64.
2. Lee AH, Ellis IO. The Nottingham prognostic index for invasive carcinoma of the breast. *Pathol Oncol Res* 2008; **14**: 113–5.
3. Tamoxifen for early breast cancer: an overview of the randomised trials. Early Breast Cancer Trialists' Collaborative Group. *Lancet* 1998; **351**: 1451–67.
4. Harbeck N, Jakesz R. St Gallen 2007: breast cancer treatment consensus report. *Breast Care* 2007; **2**: 130–4.
5. Herceptest™ Interpretation guides. available from: [http://www.dako.com/us/29036\\_herceptest\\_interpretation\\_manual-breast\\_ihc\\_us.pdf](http://www.dako.com/us/29036_herceptest_interpretation_manual-breast_ihc_us.pdf)
6. Wolff AC, Hammond ME, Schwartz JN, Hagerty KL, Allred DC, Cote RJ et al. American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol* 2007; **25**: 118–45.
7. Chin SN, Green C, Gordon-Strachan G, Wharfe G. The clinicopathologic characteristics of breast cancer at an urban academic hospital in Kingston, Jamaica. *Asian Pac J Cancer Prev* 2014; **15**: 3319–22.
8. Alfred R, Chin SN, Williams E, Walters C, Barton EN, Shah D. The prevalence and significance of oestrogen receptor (ER) positivity in breast cancer at the University Hospital of the West Indies, Jamaica. *West Indian Med J* 2012; **61**: 795–801.
9. Dent R, Trudeau M, Pritchard KI, Hanna WM, Kahn HK, Sawka CA et al. Triple-negative breast cancer: clinical features and patterns of recurrence. *Clin Cancer Res* 2007; **13**: 4429–34.
10. Carey LA, Perou CM, Livasy CA, Dressler LG, Cowan D, Conway K et al. Race, breast cancer subtypes, and survival in the Carolina Breast Cancer Study. *JAMA* 2006; **295**: 2492–502.
11. Stead LA, Lash TL, Sobieraj JE, Chi DD, Westrup JL, Charlot M et al. Triple-negative breast cancers are increased in black women regardless of age or body mass index. *Breast Cancer Res* 2009; **11**: R18. doi: 10.1186/bcr2242. [Epub 2009, Mar 25]
12. Bird PA, Hill AG, Houssami N. Poor hormone receptor expression in East African breast cancer: evidence of a biologically different disease? *Ann Surg Oncol* 2008; **15**: 1983–8.
13. Gradishar WJ, Anderson BO, Balassanian R, Blair SL, Burstein HJ, Cyr A et al. Breast Cancer Version 2.2015. *J Natl Compr Canc Netw* 2015; **13**: 448–75.
14. Seshadri R, Firgaira FA, Horsfall DJ, McCaul K, Setlur V, Kitchen P. Clinical significance of HER-2/neu oncogene amplification in primary breast cancer. The South Australian Breast Cancer Study Group. *J Clin Oncol* 1993; **11**: 1936–42.
15. Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. Human breast cancer: correlation of relapse and survival with amplification of the HER2-2/neu oncogene. *Science* 1987; **235**: 177–82.
16. Slamon DJ, Leyland-Jones B, Shak S, Fuchs H, Paton V, Bajamonde A et al. Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med* 2001; **344**: 783–92.
17. Romond EH, Perez EA, Bryant J, Suman VJ, Geyer CE Jr, Davidson NE et al. Trastuzumab plus adjuvant chemotherapy for operable HER2-positive breast cancer. *N Engl J Med* 2005; **353**: 1673–84.

© West Indian Medical Journal 2023.

This is an article published in open access under a Creative Commons Attribution International licence (CC BY). For more information, please visit [https://creativecommons.org/licenses/by/4.0/deed.en\\_US](https://creativecommons.org/licenses/by/4.0/deed.en_US).

