

## Mediastinal Thymolipoma: An Evaluation of the Clinical and Radiological Features and the Surgical Outcomes in 11 Cases

Ö Oruç<sup>1</sup>, GH Ekinci<sup>1</sup>, H Yılmaz<sup>2</sup>, A Ersev<sup>3</sup>, O Hacıomeroglu<sup>1</sup>, C Atinkaya<sup>2</sup>, T Morali<sup>1</sup>, A Yılmaz<sup>1</sup>

### ABSTRACT

**Background:** Thymolipoma is a rare benign tumour of the mediastinum, accounting for 2%–9% of all thymic tumours. Although many case reports have been published in the literature, few studies have analysed the clinical and radiological features and the surgical outcomes of this tumour.

**Objective:** To evaluate the clinical and radiological features and the surgical outcomes of the patients with thymolipoma.

**Methods:** We reviewed the records of the Pathology Department from the beginning of 2005 to the end of 2013.

**Results:** We identified 11 patients with thymolipoma. There were eight male and three female patients. Their ages ranged from 27 to 72 years, with the mean age of 40 years. All patients described pulmonary or extrapulmonary symptoms. Two patients (18.2%) had myasthenia gravis. Chest X-ray was normal in four patients. Computed tomography of the thorax revealed a mass located in the anterior mediastinum in all patients. It showed fat attenuation in 4 of 11 patients (36.4%). Thymectomy was performed in all patients. The surgical approach was thoracotomy in five, sternotomy in four and video-assisted thoracic surgery in two patients. Thymolipomas ranged in size from 4 to 33 cm. One patient died 2 years after surgery. None of the remaining patients had evidence of recurrence on follow-up.

**Conclusion:** Thymolipoma is a rare tumour of the thymus. It may be associated with myasthenia gravis. Surgical resection is the treatment of choice in the patients with thymolipoma. Complete surgical resection is the cure in most patients.

**Keywords:** Mediastinal tumour, myasthenia gravis, surgery, thymolipoma.

### INTRODUCTION

Thymolipoma is an uncommon benign tumour of the thymus consisting of mature adipose tissue and thymic tissue in variable portions (1). The term thymolipoma was first introduced in 1948 by Hall to describe this tumour (2). Up to 2008, less than 200 cases of thymolipoma had been reported in the world literature (3). This tumour occurs more frequently in young adults (4). Many patients are asymptomatic. Others may present with non-specific symptoms such as chest pain and

dyspnoea. The tumour is sometimes associated with parathymic disorders such as *myasthenia gravis*, Graves' disease, hypogammaglobulinaemia, aplastic anaemia, red cell aplasia, and Hodgkin's disease (1, 3–6). Chest X-ray may be normal or it may show an abnormal radiological appearance such as cardiomegaly, a mediastinal mass or mediastinal widening (3, 6, 7). Surgical resection is the treatment of choice in the patients with thymolipoma (5). Few studies have analysed the clinical and radiological features and the surgical outcomes of

From: <sup>1</sup>Department of Pulmonology, Süreyyapaşa Chest Diseases and Thoracic Surgery Training and Research Hospital, Istanbul, Turkey, <sup>2</sup>Department of Thoracic Surgery, Süreyyapaşa Chest Diseases and Thoracic Surgery Training and Research Hospital, Istanbul, Turkey and <sup>3</sup>Department of Pathology, Süreyyapaşa Chest Diseases and Thoracic Surgery Training and Research Hospital, Istanbul, Turkey.

Correspondence: Dr GH Ekinci, Süreyyapaşa Göğüs Hastalıkları ve Göğüs, Cerrahisi Eğitim ve Araştırma Hastanesi, Maltepe, Istanbul 34844, Turkey. Email: gulbanuh@hotmail.com

this tumour (4, 5, 7, 8). We present an evaluation of the clinical and radiological features and the surgical outcomes in 11 cases of thymolipoma.

## MATERIALS AND METHODS

This retrospective study was conducted at Süreyyapaşa Chest Diseases and Thoracic Surgery Training and Research Hospital in Istanbul, Turkey. The study was approved by the scientific committee of our institute. We reviewed pathology laboratory records from the beginning of 2005 to the end of 2013. We identified 11 cases of thymolipoma during this period. The histology of all resected tumour specimens was re-examined by a pathologist experienced in thoracic pathology. The diagnosis of thymolipoma was based on the presence of a mixture of benign thymic tissue and mature adipose tissue in the tumour at microscopic examination.

The clinical files of the patients were analysed retrospectively. For each patient, the following information was noted: age, sex, smoking habit, symptoms, radiological features, diagnostic investigations, pathological findings, treatment, and outcomes. All patients underwent routine laboratory studies and electrocardiography. Chest X-ray and computed tomography (CT) of the thorax were evaluated in all patients. Two patients were additionally evaluated with magnetic resonance imaging (MRI). 18-fluoro-deoxyglucose positron emission tomography-CT (FDG PET-CT) scan was performed in three patients. The patients' follow-up was completed as of December 2014. The hospital electronic database was reviewed to obtain follow-up information. Patients were asked about their symptoms and chest X-rays were performed, and if necessary, CT of the thorax. The modified Osserman's classification was used to assess the severity of *myasthenia gravis* throughout the patients' pre-operative care and post-operative follow-up (4).

## RESULTS

There were eight male and three female patients, indicating a male to female ratio of 2.7:1. Their ages ranged from 27 to 72 years, with the mean age of 40 years. Seven patients (63.6 %) were younger than 40 years. All patients described pulmonary and/or extrapulmonary symptoms. The most frequent symptom was chest pain. Two patients (18.2%) had *myasthenia gravis*. The severity score of the *myasthenia gravis* was IIa for one patient. She received high-dose corticosteroid therapy and pyridostigmine bromide. She had also Graves' disease. The severity score was I for the other patient. He received only pyridostigmine bromide. Table 1 shows the clinical data of the patients.

Radiological features are summarized in Table 2. Chest X-ray was normal in four patients (36.4%). Computed tomography of the thorax revealed a mass located in the anterior mediastinum in all patients. It showed fat attenuation within 4 of 11 masses (36.4%). Magnetic resonance imaging scans, obtained in two patients, demonstrated high signal intensity intermixed with areas of intermediate intensity. 18-Fluoro-deoxyglucose positron emission tomography CT scans were obtained in three patients. The maximum standardized uptake values (SUVmax) of the lesions on FDG PET-CT were 2.9, 3.2 and 3.8.

The data of treatment, pathological tumour size and outcomes of the patients are given in Table 3. All patients underwent a surgical procedure. The surgical approach was thoracotomy in five patients, sternotomy in four patients and video-assisted thoracoscopic surgery in two patients. The tumour was located in the anterior mediastinum in all cases. Only the thymus was removed in all patients. On macroscopic examination, all masses were encapsulated and soft. The tumours varied in size from 4 to 33 cm in greatest dimension, with the mean size

Table 1: The clinical data on the patients

Case	Age (years)	Gender	Smoking	Symptoms	<i>Myasthenia gravis</i>	Osserman score
1	27	Female	Yes	Cough	No	
2	34	Male	No	Cough, chest pain	No	
3	43	Male	Yes	Chest pain	No	
4	35	Female	No	Dyspnoea, weakness, Graves' disease	Yes	IIa
5	30	Female	No	Dyspnoea	No	
6	71	Male	No	Chest pain	No	
7	20	Male	No	Chest pain	No	
8	72	Male	No	Dyspnoea	No	
9	23	Male	No	Cough, chest pain, dyspnoea	No	
10	29	Male	No	Chest pain	No	
11	56	Male	Yes	Ocular findings	Yes	I

of 9 cm. Microscopic examination demonstrated a mixture of benign thymic tissue and mature adipose tissue in all tumours. No histological evidence of malignancy was seen in any of the tumours (Figure). The hospital stay ranged from 3 to 12 days, on average 6.1 days. Hematoma was developed in one patient on the 1<sup>st</sup> post-operative day following the thoracotomy and repeated thoracotomy was necessary. Another patient developed sepsis on the second post-operative day following the thoracotomy. He was treated with early aggressive antibiotic therapy and mechanical ventilatory support in an intensive care unit. There were no complications as a result of the resection of the thymolipoma in any other patients. One patient died 2 years after operation due to cardiovascular disease. None of the remaining 10 patients had evidence of recurrence or residual tumour on follow-up. The patient with severity score I

experienced a non-improvement in *myasthenia gravis* symptoms after thymectomy. The patient with severity score IIa experienced an improvement in *myasthenia gravis* symptoms after resection of thymolipoma.

## DISCUSSION

Thymolipoma is an uncommon benign tumour of the mediastinum. This tumour accounts for 2% to 9% of all thymus tumours and 1.1% of all solid mediastinal tumours (6, 9). Its incidence is approximately 0.12 cases per 100 000 inhabitants per year (10). Up to 2008, less than 200 cases of thymolipoma had been reported in the world literature (3). The pathogenesis of this tumour is unclear, but several theories have been proposed. The most widely accepted theory proposes that diffuse thymic hyperplasia is replaced by adipose tissue, the same process that occurs in the normal thymus (9, 11).

Table 2: Radiological features of the patients

CT features					
Case	Location	Site	Border	Fat on CT	
1	Widening	Anterior mediastinum	Middle	Smooth	Yes
2	Widening	Anterior mediastinum	Middle	Smooth	No
3	Widening	Anterior mediastinum	Middle	Smooth	Yes
4	Normal	Anterior mediastinum	Left	Smooth	No
5	Normal	Anterior mediastinum	Middle	Smooth	No
6	Widening	Anterior mediastinum	Left	Lobulated	No
7	Widening	Anterior mediastinum	Middle	Smooth	No
8	Widening	Anterior mediastinum	Right	Smooth	Yes
9	Mass	Anterior mediastinum	Left	Smooth	Yes
10	Normal	Anterior mediastinum	Middle	Lobulated	No
11	Normal	Anterior mediastinum	Middle	Smooth	No

CT = computed tomography.




Figure: Microscopic appearance. The tumour shows thymic tissue containing a Hassall's corpuscle admixed with adipose tissue.

Table 3: The data of treatment, pathological tumour size and outcomes of the patients

Case	Surgical approach	Operation	Hospital stay (day)	Complication	Tumour size (cm)	Follow-up (months)	Prognosis
1	Sternotomy	Thymectomy	3	No	9	44	Alive
2	Thoracotomy	Thymectomy	7	No	4.5	17	Alive
3	Thoracotomy	Thymectomy	11	No	4.5	84	Alive
4	Sternotomy	Thymectomy	5	No	9	108	Alive
5	VATS*	Thymectomy	4	No	6	15	Alive
6	Thoracotomy	Thymectomy	4	No	10	98	Alive
7	Sternotomy	Thymectomy	5	No	4	104	Alive
8	Thoracotomy	Thymectomy	12	Sepsis	8	24	Died
9	Thoracotomy	Thymectomy	11	Haematoma	33	74	Alive
10	Sternotomy	Thymectomy	3	No	4	78	Alive
11	VATS	Thymectomy	3	No	7	23	Alive

\*VATS: video-assisted thoracoscopic surgery.

Thymolipoma can occur at any age, but is more common in young adults. The majority of thymolipomas are diagnosed in the first four decades of life (7, 8, 12). There is no sex predilection (7, 9, 12). Many patients are symptom free at the time of diagnosis and the tumours are identified incidentally by chest X-ray or CT of the thorax during a diagnostic work-up for other medical problems. Others present with non-specific symptoms such as chest pain, cough, dyspnoea, respiratory tract infections, haemoptysis, dysphagia, hoarseness, tachypnoea, cyanosis, and weight loss. Most symptoms are related to the local growth of the tumour and mechanical compression of adjacent mediastinal organs (1, 5, 7–9, 11, 13). Thymolipomas often become large and they can reach a huge size (3, 5, 6, 8, 9). According to our review of the literature, the largest size of thymolipomas that has been reported in the literature is 36 cm and their weight ranges from 31 g to 16 kg (5, 8, 14). These tumours are sometimes associated with parathyroid disorders such as *myasthenia gravis*, Graves' disease, hypogammaglobulinaemia, aplastic anaemia, red cell aplasia, and Hodgkin's disease (1, 3–6). The incidence of *myasthenia gravis* associated with thymolipoma is between 2.8% and 50% (5). In the present series, all patients were symptomatic. Two patients had *myasthenia gravis*. One of them had also Graves' disease.

Radiographic features are variable. Chest X-ray may reveal an enlarged cardiac silhouette mimicking cardiomegaly. It may show a mediastinal mass or widening. The tumour may present with normal chest X-ray, as happened in our four cases (6, 7, 11, 12). Computed tomography and MRI may be helpful in the diagnosis of thymolipoma. Computed tomography reveals a mass in the anterior mediastinum. It may demonstrate fat tissue within the tumour, as shown in our four patients (3, 7, 11). Although the diagnosis of thymolipoma is strongly suggested by CT and MRI, histological examination requires to make a definitive diagnosis or even to differentiate benign from malignant tumours (3, 5, 11). The differential diagnosis of this tumour includes lipoma, mediastinal lipomatosis, mediastinal cysts, thymoma, thymic hyperplasia, teratoma, aneurysm of aorta, liposarcoma, lymphoma, and thymic carcinoma (9).

The treatment of choice is surgical resection of the tumour, which can be accomplished via sternotomy, thoracotomy or thoracoscopic approach (3, 5, 7, 11, 15). The surgical approach is chosen according to the location and extent of the tumour (3, 5). Complete surgical excision is generally curative (3, 7, 11). The tumour does not recur after surgical resection (5, 7). Surgical resection of the tumour may be followed by an improvement of myasthenic symptoms in patients with *myasthenia*

*gravis* (7). However, myasthenic symptoms may persist in some patients (5). In the present series, the patients showed no evidence of recurrence or residual tumour during the follow-up. While one patient experienced an improvement in *myasthenia gravis* symptoms after a resection of thymolipoma, myasthenic symptoms persisted in the other patient.

In conclusion, thymolipoma is a rare tumour of the thymus. It may be associated with *myasthenia gravis*. Surgical resection is the treatment of choice in the patients with thymolipoma. Complete surgical resection is curative in most patients.

## REFERENCES

1. Aghajanzadeh M, Alavi A, Pourrasouli Z, Aghajanzadeh G, Massahnia S. Giant mediastinal thymolipoma in 35 year-old women. *J Cardiovasc Thorac Res* 2011; **3**: 67–70.
2. Hall GFM. A case of thymolipoma with observations on a possible relationship to intrathoracic lipomata. *Br J Surg* 1948; **36**: 321–4.
3. Mourad OMA, de Andrade FM, Abrahao P, Monnerat A, Judice LF. Asymptomatic giant mediastinal mass: a rare case of thymolipoma. *J Bras Pneumol* 2009; **35**: 1049–52.
4. Huang CS, Li WY, Lee PC, Kao KP, Chou TY, Wu MH et al. Analysis of outcomes following surgical treatment of thymolipomatous myasthenia gravis: comparison with thymomatous and non-thymomatous myasthenia gravis. *Interact Cardiovasc Thorac Surg* 2014; **18**: 475–81.
5. Rieker RJ, Schirmacher P, Schnabel PA, Moser K, Hoffmann H, Dienemann H et al. Thymolipoma. A report of nine cases, with emphasis on its association with myasthenia gravis. *Surg Today* 2010; **40**: 132–6.
6. Tsukioka T, Inoue K, Iwata T, Mizuguchi S, Morita R, Suehiro S. Thymolipoma associated with myasthenia gravis. *General Thorac Cardiovasc Surg* 2007; **55**: 26–8.
7. Damadoglu E, Salturk C, Takir HB, Ertugrul M, Yilmaz A, Atasalihi A et al. Mediastinal thymolipoma: an analysis of 10 cases. *Respirology* 2007; **12**: 924–7.
8. Moran CA, Rosado-de-Christenson M, Suster S. Thymolipoma: clinicopathologic review of 33 cases. *Mod Pathol* 1995; **8**: 741–4.
9. Ghanem MA, Habambo G, Bahous J, Chouairy C, Khalil BA. Large thymolipoma causing primary respiratory distress. *J Med Liban* 2012; **60**: 106–9.
10. Gamondes JP, Balawi A, Greenland T. Seventeen years of surgical treatment of thymoma: factors influencing survival. *Eur J Cardiothorac Surg* 1991; **5**: 124–31.
11. Roque C, Rodriguez P, Quintero C, Santana N, Hussein M, Freixinet J. Giant thymolipoma. *Arch Bronchoneumol* 2005; **41**: 402–3.
12. Rosado-de-Christenson ML, Pugatch RD, Moran CA, Galobardes J. Thymolipoma: analysis of 27 cases. *Radiology* 1994; **193**: 121–6.
13. Hayashi A, Takamori S, Tayama K, Mitsuoka M, Ohtsuka S, Aoyama Y et al. Thymolipoma: clinical and pathological features – report of three cases and review of literature. *Kurume Med J* 1997; **44**: 141–6.
14. Kilic D, Giray S, Bolat FA, Tercan F, Bilen A, Hatipoglu A. A rare combination of thymic tumor: radiologically invisible thymolipoma associated with myasthenia gravis. *Neuro India* 2006; **54**: 322–3.
15. Tan TW, Kim DS, Wallach MT, Mangray S, Luks FI. Thoracoscopic resection of a giant thymolipoma in a 4-year-old girl. *J Laparoendosc Adv Surg Tech A* 2008; **18**: 903–5.

© West Indian Medical Journal 2021.

This is an article published in open access under a Creative Commons Attribution International licence (CC BY). For more information, please visit [https://creativecommons.org/licenses/by/4.0/deed.en\\_US](https://creativecommons.org/licenses/by/4.0/deed.en_US).

