

Chronic Polyarticular Tophaceous Gout Masquerading as Rheumatoid Arthritis

The Editor,

Sir,

Gouty arthritis most often begins as monoarticular arthritis in a lower extremity, predominantly the first metatarsophalangeal joint; however, if left untreated polyarticular involvement and development of tophi becomes more and more frequent as time passes by, especially during the chronic gouty arthritis stage (1, 2). In this stage, due to the presence of polyarthritis and subcutaneous nodules, which are typically seen in rheumatoid arthritis (RA), chronic polyarticular tophaceous gout may masquerade as RA (2, 3). Herein, we present an unusual case of chronic polyarticular gout masquerading as RA.

A 51-year-old male presented with swelling and pain of both wrists, left knee and right ankle for 3 days. He reported similar episodes over the 10 years in hands, shoulders and feet, each lasting several weeks and was alleviated with non-steroidal anti-inflammatory drugs (NSAIDs), which were his only medications. Physical examination revealed swelling and tenderness in both wrists and right ankle, and suprapatellar joint effusion in left knee. A skin coloured subcutaneous 1 × 1 cm firm nodule was seen over right achilles tendon, posteromedial to the right lateral malleolus (Fig. 1).



Fig. 1: Skin coloured subcutaneous nodules on right achilles tendon, posteromedial to the right lateral malleolus (arrows).

Symmetrical wrist involvement and presence of nodules initially led us to diagnose him as seronegative RA. Initial blood tests results were as follows: white blood cell (WBC) count: 11 400 cells/mcL, haemoglobin: 15.7 g/dL, platelet count: 245 000 cells/mcL, creatinine: 0.9 mg/dL, alanine transaminase: 53 U/L, aspartate

transaminase: 31U/L, erythrocyte sedimentation rate: 20 mm/h, and C-reactive protein: 12.1 mg/dL (N: 0–5). Anti-cyclic citrullinated peptide (CCP) and rheumatoid factor (RF) were negative. The X-rays of the hands were unremarkable, and knee radiographs showed mild degenerative changes (Fig. 2).



Fig. 2: (A) Frontal radiograph of the hands shows no significant joint narrowing and no erosive changes. (B) Frontal radiograph of the knees shows mild medial tibiofemoral joint space narrowing and minimal osteophytes.

Since RF and anti-CCP were negative, the nodules presumably were not rheumatoid and the radiographs were unremarkable even he had a 10-year history of episodes; these findings reduced the likelihood of RA. Therefore, we suspected the uncommon presentation of gouty arthritis that rarely mimics RA. His serum uric acid level was found to be 8.4 mg/dL (N: 2.5–7.5). For definitive diagnosis of gout, his left knee was aspirated. Gross appearance of synovial fluid was yellow and transparent. The WBC count was 400/mm³. Monosodium urate crystals were identified under polarized light microscopy (Fig. 3).

The patient was diagnosed as polyarticular gouty arthritis. Treatment with colchicine 1.5 mg/day was initiated for prompt and safe termination of the acute attack. After 1 month of this therapy, allopurinol 150 mg/day was added to colchicine treatment for prevention and reversal of the urate crystal depositions. Symptoms of patient had been improved, and no attack of arthritis was observed at 6-month follow-up.

Gout rarely presents with polyarthritis, subcutaneous nodules, symmetrical joint involvement and morning stiffness, which are commonly seen in RA, and the clinical presentation of these cases may mimic RA and cause diagnostic dilemma (2–5). Our patient had polyarthritis, subcutaneous nodules and symmetrical wrist involvement resembling a RA. However, subcutaneous nodules



Fig. 3: Polarized light microscopy shows negatively birefringent needle-shaped monosodium urate crystals.

are frequently seen in patients with RF-positive RA (6). Therefore, in patients with nodules and RF-negativity, other diseases including tophaceous gout should be suspected. Our patient presented with arthritis and subcutaneous nodules, and he however was seronegative for RF, which was inconsistent with RA. Furthermore, radiological images are useful to differentiate between these two diseases. Unlike the RA, radiological features of gout arthritis are asymmetrical joint involvement, preserving of joint spaces until the late stages and punched out erosions with sclerotic overhanging margins. Also, periarticular osteoporosis, which is often present in RA, is very rare in gout arthritis (7). Hand and knee graphics of our patient were unremarkable, which were also not consistent with a 10-year history of RA.

With presenting our unusual case, we would like to call attention to the rare presentation of gouty arthritis, which may masquerades as RA and so may cause a diagnostic dilemma. In clinical practice, when evaluating the patients with polyarthritis and subcutaneous nodules,

which are typical features of RA, a high index suspicion for gouty arthritis is needed.

Keywords: Gouty arthritis, polyarthritis rheumatoid arthritis, tophi.

M Karagülle¹, S Kardeş¹, P Fırat², N Erdoğan¹, MZ Karagülle¹

From: ¹Department of Medical Ecology and Hydroclimatology, Istanbul University, Istanbul, Turkey and ²Department of Pathology, Istanbul Faculty of Medicine, Istanbul University, Istanbul, Turkey.

Correspondence: Dr M Karagülle, Department of Medical Ecology and Hydroclimatology, Istanbul Faculty of Medicine, Istanbul University, Fatih, Capa, 34093 İstanbul, Turkey. Email: mkgulle@istanbul.edu.tr

REFERENCES

1. Richette P, Bardin T. Gout. *Lancet* 2010; **375**: 318–28.
2. Talbott JH, Altman RD, Yü TF. Gouty arthritis masquerading as rheumatoid arthritis or vice versa. *Semin Arthritis Rheum* 1978; **8**: 77–114.
3. Schapira D, Stahl S, Izhak OB, Balbir-Gurman A, Nahir AM. Chronic tophaceous gouty arthritis mimicking rheumatoid arthritis. *Semin Arthritis Rheum* 1999; **29**: 56–63.
4. Kotevoğlu N, Özcan B, Kuran B. Chronic tophaceous gouty arthritis resembling rheumatoid arthritis. *Arch Rheumatol* 2003; **18**: 112–5.
5. Sarmento JF, Cavalcante VDA, Sarmento MTR, Braz ADS, Freire EA. Chronic tophaceous gout mimicking rheumatoid arthritis. *Rev Bras Reumatol* 2009; **49**: 741–6.
6. Sayah A, English JC. Rheumatoid arthritis: a review of the cutaneous manifestations. *J Am Acad Dermatol* 2005; **53**: 191–209.
7. Resnick G, Niwayama G. *Diagnosis of bone and joint disorders*. 2nd ed. Philadelphia: W.B. Saunders Company; 1988.

© West Indian Medical Journal 2023.

This is an article published in open access under a Creative Commons Attribution International licence (CC BY). For more information, please visit https://creativecommons.org/licenses/by/4.0/deed.en_US.

